

# Climate-Resilient Agriculture for Tribal Farmers — Building From What Exists

Odisha has always been weather-volatile. The state experiences cyclones, droughts, floods, and erratic monsoons — sometimes in the same year. What is changing under climate change is the frequency and unpredictability of extremes:

JABASU KNOWLEDGE COMMONS · JABASU.ORG

---

**B** practice-note Agriculture & Markets

Published: April 2026 · Last reviewed: April 2026

**Monsoon onset timing:** Traditional agricultural calendars — when to sow, when to transplant, when to apply compost — are calibrated against historical monsoon patterns. Delayed or erratic monsoon onset disrupts these calendars in ways that cascades through the whole season. Farmers who sow based on historical dates into soil that isn't yet adequately moist face germination failure.

**Extended dry spells within the monsoon:** Even in years of normal total rainfall, extended dry spells within the monsoon season — two to three weeks without rain during the period when crops most need it — are increasing in frequency. Rain-fed crops without supplemental irrigation cannot bridge these gaps.

**Temperature increases:** Higher temperatures during the growing season are reducing yields of heat-sensitive crops and disrupting the timing of flowering and grain fill. For millets — traditionally the most heat and drought-tolerant of tribal staple crops — even these hardy varieties face stress at the upper temperature extremes now being recorded.

**Post-monsoon rains and harvest timing:** Late-season rains that arrive when crops should be harvested cause grain losses, fungal disease outbreaks, and storage quality problems.

**Forest fire season extension:** Longer, hotter dry seasons are extending the period when forest fires are a risk — affecting both NTFP availability and agricultural land on forest margins.

Against this intensifying climate stress, tribal farmers in Odisha's highland districts are managing with fewer resources and less institutional support than almost any comparable farming community in India.

---

# What Traditional Tribal Farming Systems Already Do Right

Before designing any climate adaptation programme for tribal farmers, programme designers need to understand what the existing farming system is already doing — and why. The following are not quaint customs. They are functional adaptations.

## Crop diversity as insurance

Most tribal farmers in highland Odisha grow multiple crops simultaneously — some combination of millets (finger millet, little millet, kodo, foxtail), pulses (horse gram, pigeon pea, cowpea), oilseeds (Niger), vegetables, and sometimes paddy where water is available. This diversity is not inefficiency. It is portfolio diversification. When the rainfall pattern favours drought-tolerant crops, the farmer has those. When rains are adequate, the higher-yielding crops do well. No single failure destroys the season.

The PMC (Pure Millet Crop) approach — which some agricultural extension programmes have promoted, replacing mixed cropping with single-crop millet production — is agronomically simpler and easier to measure, and produces lower climate resilience. The research consistently supports mixed cropping over monoculture in rain-fed systems.

## Traditional seed varieties as climate adaptation

Locally-adapted traditional seed varieties — selected by farmers over generations for performance in their specific micro-climate — carry drought tolerance, pest resistance, and nutritional characteristics that bred hybrid varieties do not. They are also free. Farmers who save and exchange their own seeds are not dependent on market supply chains that may fail at exactly the moment they are needed (drought years, when input dealers' stocks are depleted).

The PMC Frontiers in Plant Science analysis (2025) documents finger millet's exceptional resilience: it flourishes in marginal and degraded lands, withstands drought, poor soil, and temperature stress, and requires minimal water and external inputs compared to any alternative. This is not a new discovery. Tribal farmers in Odisha have grown it for this reason for generations. What they often lack is: sufficient seed stock of locally-adapted varieties; knowledge of improved agronomic practices (spacing, composting) that increase yields from the same resilient varieties; and markets that reward the crop's resilience with adequate prices.

## Traditional water harvesting and soil conservation

Check dams (bandhs), terracing, and traditional water harvesting structures built and maintained by communities in tribal highland areas are climate adaptation infrastructure that in many cases predates any government irrigation investment. Where these have been maintained, they buffer the impacts of dry spell intensification. Where they have been allowed

to deteriorate, dry spells that communities previously managed through these structures now cause crop failure.

---

## What Climate-Smart Agriculture Adds — and What It Doesn't Replace

"Climate-smart agriculture" (CSA) is a framework promoted by FAO, McKinsey, and international development bodies that describes agricultural practices that simultaneously improve food security, enhance resilience, and reduce or remove greenhouse gas emissions. In the smallholder Indian context, the most relevant CSA practices for Odisha's tribal farmers are:

**Seed bank development:** Community seed banks (covered in the Millet Value Chains Practice Note) that conserve diverse, locally-adapted seed varieties are the single most important climate adaptation infrastructure for tribal farming communities. They provide access to the variety diversity that makes crop portfolio resilience possible, and they insulate communities from the market failures in commercial seed supply that typically worsen during climate stress events.

The Sangham Seed Bank in Telangana — managed by women's groups, preserving 80 types of traditional food crops — is the benchmark model. Comparable community seed banks in Odisha's tribal districts, facilitated through the OMM or independently, are high-priority climate adaptation investments.

**Water harvesting and tank restoration:** Small-scale water harvesting structures — check dams, percolation ponds, field bunds — buffer dry spell impacts significantly. McKinsey's analysis of climate adaptation for Indian smallholders found water management interventions among the highest-return adaptations available, noting that almost every smallholder can adopt at least one water management adaptation measure, and that those who adopt multiple measures achieve greater resilience.

MGNREGS, in tribal PESA areas where gram sabhas control worksite selection, is the primary funding mechanism for community-designed water harvesting. An active gram sabha that selects water harvesting works on land adjacent to its agricultural area, rather than accepting whatever works the block office prescribes, is doing climate adaptation.

**System of Millet Intensification (SMI):** Line sowing or transplanting at wider spacing — the approach piloted under the OMM — produces 20–40% yield improvement from the same locally-adapted varieties that communities already grow. It does not require external inputs beyond the changed practice. It increases the income per hectare from climate-resilient crops, making those crops economically competitive with paddy even in years when water is adequate for paddy cultivation.

**Bio-inputs replacing chemical fertilisers:** Compost, jeevamrut, and locally-produced bio-fertilisers improve soil organic matter — which improves soil water retention, reduces the impact of dry spells, and produces crop yield benefits in the medium term. Soil with higher organic matter content experiences less yield loss during a dry spell than soil with low organic matter. The transition to bio-inputs is both a productivity intervention and a climate adaptation.

**Agroforestry integration:** Trees on or adjacent to agricultural land moderate temperature, provide organic matter through leaf fall, create microclimatic buffers, and provide alternative income during crop failure years through fruits, nuts, and timber. Traditional tribal farming systems often include tree integration; agricultural extension programmes have historically encouraged field clearance rather than agroforestry. Supporting communities to maintain and restore tree integration in their agricultural landscapes is evidence-based climate adaptation.

---

## What NGOs Should Not Do

**Promote hybrid monoculture in rain-fed tribal contexts.** The evidence across contexts is consistent: monoculture of high-yielding varieties in rain-fed systems increases income in optimal weather and collapses in bad weather. The same family that produces a bumper yield from hybrid paddy in a good monsoon year produces nothing in a drought year. The tribal farmer growing mixed millets and pulses on traditional varieties produces less in the good year and much more in the bad year. Climate adaptation favours the latter.

**Import standard "climate-smart" curricula without contextualisation.** A climate-smart agriculture training module designed for peri-urban Maharashtra farmers, or for irrigated systems in Punjab, is not climate-smart in a tribal highland in Malkangiri. The adaptation measures, the crop systems, the social structures, the risk profiles are all different. Build from the existing system, add specific technical improvements, test them with community members before scaling.

**Dismiss traditional seasonal calendars.** Tribal farmers' traditional agricultural calendars — which plants to sow after which phenological indicators, which practices to follow after which weather signs — contain accumulated observational data that formal agro-meteorology often cannot match at the local level. Climate change is disrupting some of these indicators, and communities need support to update their calendars accordingly. But the calendar-keeping practice itself is a climate resilience tool, not an obstacle to it.

---

## A Practical Programme Design Sequence

For an NGO wanting to support climate-resilient agriculture among tribal farmers in Odisha:

**First:** Conduct a participatory agricultural calendar exercise with farmers in the target community. What do they sow, when, in response to what indicators? Where is climate variability already disrupting their calendar? What practices have they adapted informally?

**Second:** Map seed diversity. What varieties of each crop does the community currently grow? What varieties have been lost in the last ten years? What varieties do elders remember that are no longer grown? This mapping is both documentation and the foundation for seed bank design.

**Third:** Establish a community seed bank using the most valued, locally-adapted varieties identified in step two. This is a one-to-two-month facilitation process that produces the core climate adaptation infrastructure.

**Fourth:** Introduce SMI (for millets) and bio-input preparation (for all crops) as field demonstrations on volunteer farmers' plots. Do not prescribe — demonstrate and let the results speak. Farmers who see their neighbour's SMI plot yielding 30% more from the same ragi seed they both saved will adopt.

**Fifth:** Facilitate gram sabha planning for MGNREGS water harvesting works. Connect the agricultural vulnerability assessment from step one to the gram sabha's worksite selection choices under MGNREGS.

This sequence — observation first, documentation second, infrastructure third, agronomy fourth, system investment fifth — builds from what communities know before adding what they might adopt. It is slower than arriving with a training module and a demonstration plot. It produces more durable outcomes.

---

Related Knowledge Commons content: Environment & Climate Sector Primer (Sector 07) · Agriculture & Markets Sector Primer (Sector 09) · Practice Note: Millet Value Chains — The OMM Model · Practice Note: Community Forest Rights — From Title to Conservation

Evidence Grade: B — Multi-study. This Practice Note draws on PMC Frontiers in Plant Science millet resilience analysis (2025), McKinsey climate-smart agriculture for smallholders analysis (2023), Cutter Consortium climate-resilient agriculture case studies, AIF climate-smart agriculture documentation, and WASSAN field documentation on System of Millet Intensification outcomes in Odisha. Last reviewed: April 2026.

Questions or corrections: [knowledge@jabasu.org](mailto:knowledge@jabasu.org)